

=> d his

(FILE 'HOME' ENTERED AT 16:04:47 ON 16 MAY 2006)

FILE 'REGISTRY' ENTERED AT 16:05:20 ON 16 MAY 2006

L1 STRUC
L2 0 S L1
L3 2 S L1 FUL
L4 STRUC
L5 3 S L4
L6 3 S L5 NOT L3

FILE 'CAPLUS' ENTERED AT 16:10:12 ON 16 MAY 2006

L7 1 S L6
L8 2 S L3
L9 41 S (DEACETYLASE(L) INHIBITOR?) AND PIPERIDIN?
L10 41 S L9 NOT (L7 OR L8)
L11 23 S L10 AND (PYRROL? OR PYRAZO? OR OXAZO? OR FURAN? OR THIEN?)

=> s l10 and benzimidaz?

32894 BENZIMIDAZ?

L12 10 L10 AND BENZIMIDAZ?

=> s l12 and l11


L13 8 L12 AND L11

=> d bib abs 1-8

L13 ANSWER 1 OF 8 CAPLUS COPYRIGHT 2006 ACS on STN
AN 2005:1075803 CAPLUS
DN 143:367317
TI Preparation of N-(2-amino and 2-hydroxy)phenyl carboxamides as
inhibitors of histone deacetylase
IN Delorme, Daniel; Vaisburg, Arkadii; Moradei, Oscar; Leit, Silvana;
Raepfel, Stephane; Frechette, Sylvie; Bouchain, Giliiane; Zhou, Zhihong;
Paquin, Isabelle; Gaudette, Frederic; Isakovic, Ljubomir
PA Methylgene Inc., Can.
SO PCT Int. Appl., 245 pp.
CODEN: PIXXD2
DT Patent
LA English
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	WO 2005092899	A1	20051006	WO 2005-CA454	20050329
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	US 2005245518	A1	20051103	US 2005-90713	20050325
PRAI	US 2004-556828P	P	20040326		
	US 2005-90713	A	20050325		
	WO 2005-IB802	A	20050325		
OS	MARPAT 143:367317				
GI					

=> d l1
 L1 HAS NO ANSWERS
 L1 STR



NODE ATTRIBUTES:
 DEFAULT MLEVEL IS ATOM
 DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:
 RSPEC 9 16
 NUMBER OF NODES IS 27

STEREO ATTRIBUTES: NONE

=> s l1 ful
 FULL SEARCH INITIATED 16:07:51 FILE 'REGISTRY'
 FULL SCREEN SEARCH COMPLETED - 41 TO ITERATE


100.0% PROCESSED 41 ITERATIONS
 SEARCH TIME: 00.00.04

2 ANSWERS

L3 2 SEA SSS FUL L1

=> d 1-2

L3 ANSWER 1 OF 2 REGISTRY COPYRIGHT 2006 ACS on STN
 RN 758693-31-1 REGISTRY
 ED Entered STN: 08 Oct 2004
 CN 2-Propenamide, N-hydroxy-3-[3-[1-(1-methyl-4-piperidiny)]-1H-benzimidazol-2-yl]phenyl]- (9CI) (CA INDEX NAME)
 FS 3D CONCORD
 MF C22 H24 N4 O2
 SR CA
 LC STN Files: CA, CAPLUS, USPATFULL





PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

1 REFERENCES IN FILE CA (1907 TO DATE)
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L3 ANSWER 2 OF 2 REGISTRY COPYRIGHT 2006 ACS on STN
RN 404949-04-8 REGISTRY
ED Entered STN: 10 Apr 2002
CN 1-Piperidinecarboxylic acid, 4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-, 2-[[[4-[(1E)-3-(hydroxyamino)-3-oxo-1-propenyl]phenyl]methyl][2-(1H-indol-3-yl)ethyl]amino]ethyl ester (9CI) (CA INDEX NAME)
FS STEREOSEARCH
MF C35 H38 N6 O5
SR CA
LC STN Files: CA, CAPLUS, TOXCENTER, USPAT2, USPATFULL

Double bond geometry as shown.





VAR G1=13/14
 ENTER (DIS), GRA, NOD, BON OR ?:end
 L4 STRUCTURE CREATED

=> s 14
 SAMPLE SEARCH INITIATED 16:09:42 FILE 'REGISTRY'
 SAMPLE SCREEN SEARCH COMPLETED - 3 TO ITERATE

100.0% PROCESSED 3 ITERATIONS 3 ANSWERS
 SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
 BATCH **COMPLETE**
 PROJECTED ITERATIONS: 3 TO 163
 PROJECTED ANSWERS: 3 TO 163


L5 3 SEA SSS SAM L4

=> s 15 not 13
 L6 3 L5 NOT L3

=> d scan

L6 3 ANSWERS REGISTRY COPYRIGHT 2006 ACS on STN
 IN 2-Propenamide, 3-[3-[1-[(3R)-1-ethyl-3-piperidinyl]-5,6-difluoro-1H-
 benzimidazol-2-yl]phenyl]-N-hydroxy- (9CI)
 MF C23 H24 F2 N4 O2

Absolute stereochemistry.
 Double bond geometry unknown.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):0

=> fil caplus

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

173.82

174.03

FILE 'CAPLUS' ENTERED AT 16:10:12 ON 16 MAY 2006

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 16 May 2006 VOL 144 ISS 21

FILE LAST UPDATED: 15 May 2006 (20060515/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/infopolicy.html>

=> s 16

L7 1 L6

=> d bib abs hitstr

L7 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2004:799454 CAPLUS

DN 141:291229

TI Histone deacetylase inhibitors

IN Bressi, Jerome C.; Brown, Jason W.; Cao, Sheldon X.; Gangloff, Anthony R.; Jennings, Andrew J.; Stafford, Jeffrey A.; Vu, Phong H.; Xiao, Xiao-Yi

PA Syrrx, Inc., USA

SO PCT Int. Appl., 276 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	---	-----	-----	-----
PI	WO 2004082638	A2	20040930	WO 2004-US8342	20040317
	WO 2004082638	A3	20050506		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW:	BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE,				

ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI,
SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN,
TD, TG

CA 2518318	AA	20040930	CA 2004-2518318	20040317
US 2004254220	A1	20041216	US 2004-803575	20040317
US 2004266769	A1	20041230	US 2004-803344	20040317
US 2005137232	A1	20050623	US 2004-803580	20040317
EP 1608628	A2	20051228	EP 2004-757631	20040317

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK

PRAI US 2003-455437P P 20030317
US 2003-531203P P 20031219
WO 2004-US8342 W 20040317

OS MARPAT 141:291229


AB Compds. that may be used to inhibit histone deacetylase are disclosed.
Thus, 119 compds. were prepared which exhibited better than 1000 nM IC50
against HDAC1, HDAC2, HDAC6, and HDAC8 (suberanilohydroxamic acid showed
an IC50 of 63 nM in this assay). Many of these compds. were
3-[3-(1-substituted-1H-benzoimidazol-2-yl)phenyl]acrylic acids and
N-hydroxy-[3-(1-substituted-1H-benzoimidazol-2-yl)phenyl]acrylamides.

IT 758693-30-0 758694-08-5 758694-10-9

RL: BSU (Biological study, unclassified); PRP (Properties); BIOL
(Biological study)
(histone deacetylase inhibitors)

RN 758693-30-0 CAPLUS


CN 2-Propenamide, 3-[3-[1-(1-ethyl-3-piperidiny)]-1H-benzimidazol-2-
yl]phenyl]-N-hydroxy- (9CI) (CA INDEX NAME)



RN 758694-08-5 CAPLUS

CN 2-Propenamide, 3-[3-[1-[(3R)-1-ethyl-3-piperidiny]]-5,6-difluoro-1H-
benzimidazol-2-yl]phenyl]-N-hydroxy- (9CI) (CA INDEX NAME)


Absolute stereochemistry.
Double bond geometry unknown.



RN 758694-10-9 CAPLUS

CN 2-Propenamide, 3-[3-[1-[(3R)-1-ethyl-3-piperidiny]]-5,6,7-trifluoro-1H-
benzimidazol-2-yl]phenyl]-N-hydroxy- (9CI) (CA INDEX NAME)

Absolute stereochemistry.
Double bond geometry unknown.



=> s 13

L8 2 L3

=> d bib abs hitstr 1-2

L8 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2006 ACS on STN
AN 2004:799454 CAPLUS
DN 141:291229
TI Histone deacetylase inhibitors
IN Bressi, Jerome C.; Brown, Jason W.; Cao, Sheldon X.; Gangloff, Anthony R.;
Jennings, Andrew J.; Stafford, Jeffrey A.; Vu, Phong H.; Xiao, Xiao-Yi
PA Syrrx, Inc., USA
SO PCT Int. Appl., 276 pp.
CODEN: PIXXD2
DT Patent
LA English
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004082638	A2	20040930	WO 2004-US8342	20040317
	WO 2004082638	A3	20050506		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
	RW:	BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
	CA 2518318	AA	20040930	CA 2004-2518318	20040317
	US 2004254220	A1	20041216	US 2004-803575	20040317
	US 2004266769	A1	20041230	US 2004-803344	20040317
	US 2005137232	A1	20050623	US 2004-803580	20040317
	EP 1608628	A2	20051228	EP 2004-757631	20040317
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK			
PRAI	US 2003-455437P	P	20030317		
	US 2003-531203P	P	20031219		
	WO 2004-US8342	W	20040317		

OS MARPAT 141:291229

AB Compds. that may be used to inhibit histone deacetylase are disclosed. Thus, 119 compds. were prepared which exhibited better than 1000 nM IC50 against HDAC1, HDAC2, HDAC6, and HDAC8 (suberanilohydroxamic acid showed an IC50 of 63 nM in this assay). Many of these compds. were


3-[3-(1-substituted-1H-benzoimidazol-2-yl)phenyl]acrylic acids and
N-hydroxy-3-[3-(1-substituted-1H-benzoimidazol-2-yl)phenyl]acrylamides.

IT 758693-31-1

RL: BSU (Biological study, unclassified); PRP (Properties); BIOL
(Biological study)
(histone deacetylase inhibitors)

RN 758693-31-1 CAPLUS

CN 2-Propenamide, N-hydroxy-3-[3-[1-(1-methyl-4-piperidiny)-1H-benzimidazol-
2-yl]phenyl]- (9CI) (CA INDEX NAME)



L8 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2006 ACS on STM

AN 2002:220554 CAPLUS

DN 136:262995

TI Preparation of hydroxamic acids as deacetylase inhibitors

IN Bair, Kenneth Walter; Green, Michael A.; Perez, Lawrence B.; Remiszewski,
Stacy W.; Sambucetti, Lidia; Versace, Richard William; Sharma, Sushil
Kumar

PA Novartis AG, Switz.; Novartis-Erfindungen Verwaltungsgesellschaft mbH;
Novartis Pharma GmbH

SO PCT Int. Appl., 96 pp.

CODEN: PIXXD2



DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002022577	A2	20020321	WO 2001-EP10037	20010830
	WO 2002022577	A3	20020906		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
	CA 2420899	AA	20020321	CA 2001-2420899	20010830
	AU 2001082129	A5	20020326	AU 2001-82129	20010830
	BR 2001013669	A	20030603	BR 2001-13669	20010830
	EP 1318980	A2	20030618	EP 2001-960717	20010830
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
	JP 2004509105	T2	20040325	JP 2002-526830	20010830
	NZ 524365	A	20041126	NZ 2001-524365	20010830
	US 2003018062	A1	20030123	US 2001-944275	20010831
	US 6552065	B2	20030422		
	US 2004024067	A1	20040205	US 2002-299518	20021116
	ZA 2003001423	A	20040421	ZA 2003-1423	20030221

	NO 2003000867	A	20030225	NO 2003-867	20030225
	US 2005085507	A1	20050421	US 2004-984501	20041109
PRAI	US 2000-229943P	P	20000901		
	US 2001-292232P	P	20010518		
	US 2001-307490P	P	20010724		
	WO 2001-EP10037	W	20010830		
	US 2001-944275	A1	20010831		
	US 2002-299518	A1	20021116		
OS	MARPAT 136:262995				
GI					



II

AB The title compds. [I; R1 = H, halo, alkyl; R2 = H, alkyl, cycloalkyl, etc.; R3, R4 = H, alkyl, acyl, acylamino; or R3 and R4 together with the carbon atom to which they are bound = CO, CS, C:NR8; or R2 together with the N atom to which is bound and R3 together with the C atom to which it is bound form heterocycloalkyl, heteroaryl, etc.; R5 = H, alkyl, aryl, etc.; n1-n3 = 0-6; X, Y = H, halo, alkyl, etc.; R8 = H, alkyl, aryl, etc.] which are deacetylase inhibitors and therefore suitable for pharmaceutical compns. having anti-proliferative properties, were prepared E.g., a 3-step synthesis of II, starting with 4-formylcinnamic acid, was given. The exemplified compds. I showed IC50 of 0.005-0.5 μ M against HDA.

IT 404949-04-8P


RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of hydroxamic acids as deacetylase inhibitors)

RN 404949-04-8 CAPLUS

CN 1-Piperidinecarboxylic acid, 4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-, 2-[[[4-[(1E)-3-(hydroxyamino)-3-oxo-1-propenyl]phenyl]methyl][2-(1H-indol-3-yl)ethyl]amino]ethyl ester (9CI) (CA INDEX NAME)

Double bond geometry as shown.



=> s (deacetylase(l)inhibitor?) and piperidin?

5619 DEACETYLASE


982475 INHIBITOR?

2646 DEACETYLASE (L) INHIBITOR?

92742 PIPERIDIN?

L9

41 (DEACETYLASE (L) INHIBITOR?) AND PIPERIDIN?



AB The invention relates to N-(2-amino and 2-hydroxy)phenyl carboxamides (2-TC₆H₄NHC(O)(CH:CH)qAr-X-Cy (I); variables defined below; e.g. (E)-N-(2-Aminophenyl)-3-[4-[[2-(2-hydroxyethyl)[2-(1H-indol-3-yl)ethyl]amino]methyl]phenyl]acrylamide (shown as II)) useful for inhibiting histone **deacetylase** (HDAC) enzymic activity. The invention also provides a method for inhibiting histone **deacetylase** in a cell using said compds. as well as a method for treating cell proliferative diseases and conditions using said HDAC inhibitors. Further, the invention provides pharmaceutical compns. comprising the HDAC inhibiting compds. and a pharmaceutically acceptable carrier. For I: Cy is aryl, heteroaryl, cycloalkyl, or heterocyclyl, each of which is (un)substituted and each of which is optionally fused to ≥1 aryl or heteroaryl rings, or to ≥1 saturated or partially unsatd. cycloalkyl or heterocyclic rings, each of which rings is (un)substituted; X = a chemical bond, L, W-L, L-W, and L-W-L, wherein W, at each occurrence, is S, O, C:O, or N(R₉), where R₉ = H, alkyl, hydroxyalkyl, and tert-butoxycarbonyl; and L = C1-C4 alkylene; Ar is arylene or heteroarylene, each of which is (un)substituted; q = 0-1; and T is NH₂ or OH, provided that when Cy is naphthyl, X is -CH₂-, Ar is Ph, and q = 0-1, T is not OH. Although the methods of preparation are not claimed, 215 example preps. and/or characterization data are included. For example, II was prepared in 6 steps (59, 83, 97, 79, 96 and 80 % yields) starting from (E)-4-formylcinnamic acid and involving intermediates Me (E)-3-(4-formylphenyl)acrylate, Me (E)-3-[4-[[2-(1H-indol-3-yl)ethyl]amino]methyl]phenyl]acrylate, Me (E)-3-[4-[[2-[(tert-butyl)dimethylsilyl]oxy]ethyl][2-(1H-indol-3-yl)ethyl]amino]methyl]phenyl]acrylate, (E)-3-[4-[[2-[(tert-butyl)dimethylsilyl]oxy]ethyl][2-(1H-indol-3-yl)ethyl]amino]methyl]phenyl]acrylic acid and (E)-N-(2-aminophenyl)-3-[4-[[2-[(tert-butyl)dimethylsilyl]oxy]ethyl][2-(1H-indol-3-yl)ethyl]amino]methyl]phenyl]acrylamide.

RE.CNT 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 2 OF 8 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2005:300395 CAPLUS

DN 142:355054

TI Preparation of amide derivatives as inhibitors of histone **deacetylase**

IN Moradei, Oscar; Paquin, Isabelle; Leit, Silvana; Frechette, Sylvie; Vaisburg, Arkadii; Besterman, Jeffrey M.; Tessier, Pierre; Mallais, Tammy C.

PA Methylgene, Inc., Can.


SO PCT Int. Appl., 559 pp.
CODEN: PIXXD2

DT Patent


LA English

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2005030705	A1	20050407	WO 2004-US31591	20040924
	WO 2005030705	C2	20060420		
	W:				
	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW,				
	RW:				
	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
PRAI	US 2003-505884P	P	20030924		
	US 2003-532973P	P	20031229		
	US 2004-561082P	P	20040409		
OS	MARPAT 142:355054				
GI					



I



II

AB Title compds. I [Ar1 = (un)saturated-, (un)substituted-mono or fused poly-cyclic hydrocarbyl optionally containing 1-4 heteroatoms per ring; R1 = (un)substituted-mono-, -bi-, -tri-cyclic-aryl or -heteroaryl; R2, R3, and R4 independently = H, halo, amino, etc.; R5 and R6 independently = H, alkyl, aryl, etc.; x = 0-1; Y = any pharmaceutically acceptable chemical moiety consisting of 1 to 50 atoms with provisions] and their pharmaceutically acceptable salts, are prepared and disclosed as inhibitors of histone deacetylase. Thus, e.g., II was prepared by Suzuki coupling of 2-bromo-2-nitro-phenylamine (preparation given) with 2-thiopheneboronic acid followed by carbonylation with

4-[3,4-dimethoxy-(phenylamino)-methyl]benzoic acid (preparation given) and subsequent reduction The inhibitory capability of I towards antiproliferative activity of histone deacetylase enzyme was evaluated using 3-[4,5-dimethylthiazol-2-yl-2,5-diphenyltetrazolium] bromide (MTT) assay and it revealed that certain compds. of the invention had MTT IC 50 values in the range of below 1 up to 20 μ M. I as histone deacetylase inhibitors should prove useful in the treatment of diseases such as, but not limited to, cell proliferative disease, protozoal disease, and fungal disease.

RE.CNT 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 3 OF 8 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2005:300394 CAPLUS

DN 142:373563

TI Preparation of amide derivatives as inhibitors of histone deacetylase

IN Moradei, Oscar; Paquin, Isabelle; Leit, Silvana; Frechette, Sylvie; Vaisburg, Arkadii; Besterman, Jeffrey M.; Tessier, Pierre; Mallais, Tammy C.

PA Methylgene, Inc., Can.

SO PCT Int. Appl., 389 pp.
CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 2

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005030704	A1	20050407	WO 2004-US31590	20040924
W:				
AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW:				
BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				



PRAI US 2003-505884P P 20030924

US 2003-532973P P 20031229

US 2004-561082P P 20040409

OS MARPAT 142:373563

GI




AB Title compds. I [Arl = (un)saturated-, (un)substituted-mono or fused poly-cyclic hydrocarbonyl optionally containing 1-4 heteroatoms per ring; R1 = (un)substituted-mono-, -bi-, -tri-cyclic-aryl or -heteroaryl; R2, R3, and R4 independently = H, halo, amino, etc.; R5 and R6 independently = H, alkyl, aryl, etc.; x = 0-1; Y = any pharmaceutically acceptable chemical moiety consisting of 1 to 50 atoms with provisions] and their pharmaceutically acceptable salts, are prepared and disclosed as inhibitors of histone deacetylase. Thus, e.g., II was prepared by Suzuki coupling of 2-bromo-2-nitro-phenylamine (preparation given) with 2-thiopheneboronic acid followed by carbonylation with 4-[3,4-dimethoxy-(phenylamino)-methyl]benzoic acid (preparation given) and subsequent reduction. The inhibitory capability of I towards antiproliferative activity of histone deacetylase enzyme was evaluated using 3-[4,5-dimethylthiazol-2-yl-2,5-diphenyltetrazolium] bromide (MTT) assay and it revealed that certain compds. of the invention had MTT IC 50 values in the range of below 1 up to 20 μ M. I as histone deacetylase inhibitors should prove useful in the treatment of diseases such as, but not limited to, cell proliferative disease, protozoal disease, and fungal disease.

RE.CNT 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 4 OF 8 CAPLUS COPYRIGHT 2006 ACS on STN
AN 2004:589250 CAPLUS
DN 141:140470
TI Preparation of aminophenylbenzamides as inhibitors of histone deacetylase
IN Delorme, Daniel; Zhou, Zhihong
PA Methygene, Inc., Can.
SO U.S. Pat. Appl. Publ., 318 pp., Cont.-in-part of U.S. Ser. No. 242,304.
CODEN: USXXCO
DT Patent
LA English
FAN.CNT 3
PATENT NO. KIND DATE APPLICATION NO. DATE

PI	US 2004142953	A1	20040722	US 2003-358556	20030204
	US 6897220	B2	20050524		
	US 2004106599	A1	20040603	US 2002-242304	20020912
	AU 2004210016	A1	20040819	AU 2004-210016	20040204
	CA 2515338	AA	20040819	CA 2004-2515338	20040204
	WO 2004069823	A1	20040819	WO 2004-CA139	20040204
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	EP 1590340	A1	20051102	EP 2004-707852	20040204
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
	CN 1723207	A	20060118	CN 2004-80001769	20040204
	BR 2004007195	A	20060214	BR 2004-7195	20040204
	US 2006058298	A1	20060316	US 2005-81095	20050315
	JP 2005255683	A2	20050922	JP 2005-80310	20050318
	US 2005288282	A1	20051229	US 2005-91025	20050325
PRAI	US 2001-322402P	P	20010914		
	US 2002-391728P	P	20020626		
	US 2002-242304	A2	20020912		
	JP 2003-528544	A3	20020912		
	US 2003-358556	A	20030204		
	WO 2004-CA139	W	20040204		
OS	MARPAT 141:140470				
GI					



AB Title compds. e.g. (I; Y, Z = N, CH; W = Q1, Q2, Q3, etc.), were prepared Thus, 4-[[[4-Amino-6-(2-indanylamino)-[1,3,5]triazin-2-yl)amino]methyl]benzoic acid (preparation given) in DMF was stirred with Et₃N, BOP, and 1,2-phenylenediamine to give 63% 4-[[[4-Amino-6-(2-indanylamino)-[1,3,5]triazin-2-yl)amino]methyl]-N-(2-aminophenyl)benzamide. The latter inhibited human histone deacetylase HDAC-1 with IC₅₀ = 0.4 μM.

RE.CNT 27 THERE ARE 27 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 5 OF 8 CAPLUS COPYRIGHT 2006 ACS on STN
AN 2003:633649 CAPLUS
DN 139:179896
TI Preparation of biphenyl hydroxamic acids as inhibitors of histone deacetylase useful against cancer
IN Leahy, Ellen M.; Verner, Erik J.
PA Axys Pharmaceuticals, USA

SO PCT Int. Appl., 135 pp.


CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003066579	A2	20030814	WO 2003-US3846	20030207
	WO 2003066579	A3	20031030		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
	CA 2473505	AA	20030814	CA 2003-2473505	20030207
	AU 2003215112	A1	20030902	AU 2003-215112	20030207
	US 2004091951	A1	20040513	US 2003-360534	20030207
	EP 1472216	A2	20041103	EP 2003-710929	20030207
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK			
	JP 2005517007	T2	20050609	JP 2003-565954	20030207
	US 2006058553	A1	20060316	US 2005-503508	20051012
PRAI	US 2002-355700P	P	20020207		
	WO 2003-US3846	W	20030207		
OS	MARPAT 139:179896				
GI					




I

AB The present invention is directed to certain bicyclic hydroxamic acids (shown as I; variables defined below; e.g. N-hydroxy-4-(3-methoxyphenyl)benzamide) that are inhibitors of histone deacetylase (no data) and are therefore useful in the treatment of diseases associated with histone deacetylase activity. Pharmaceutical compns. (5 examples) and processes for preparing these compds. are also disclosed. For I: R¹ is H or alkyl; R² is H; Ar¹ is phenylene or a six membered heteroarylene ring containing one or two N ring atoms, the rest of the ring atoms being C; wherein said Ar¹ group is (un)substituted with one or two alkyl, halo, hydroxy, alkoxy, haloalkoxy, or haloalkyl; Ar² is aryl, benzimidazol-2-yl, cycloalkyl or heterocycloalkyl; R³ is H, alkyl, halo, hydroxy, or alkoxy. R⁴ and R⁵ = H, alkyl, halo, haloalkyl, nitro, cyano, carboxy, carboxyalkyl, alkoxycarbonyl, (un)substituted Ph, (un)substituted heteroaryl, (un)substituted heterocycloalkyl, cycloalkyl, heterocycloaminoalkyl, -X-R⁶, or -(C1-6alkylene)-Y-R⁷ where X and Y = -O-, -S-, -SO-, -SO₂-, -NR⁸-, -CO-, -NR⁹CO-, -CONR¹⁰-, -NR¹¹SO₂-, -SO₂NR¹²-, -NHC(O)O-, -OC(O)NH-, -NR¹³CONR¹⁴-, or -NR¹⁵SO₂NR¹⁶-; addnl. details including provisos are given in the claims. Although the methods of preparation are not claimed, .apprx.20 example prepns. of I are included.

AN 2003:242160 CAPLUS
 DN 138:271705
 TI Preparation of triazinyl and other carboxamides as inhibitors of histone deacetylase
 IN Delorme, Daniel; Woo, Soon Hyung; Vaisburg, Arkadii; Moradel, Oscar; Leit, Silvana; Raeppe, Stephane; Frechette, Sylvie; Bouchain, Giliane
 PA Methylgene, Inc., Can.
 SO PCT Int. Appl., 347 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 3

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003024448	A2	20030327	WO 2002-US29017	20020912
	WO 2003024448	A3	20031113		
	W:				
	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW				
	RW:				
	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	CA 2465978	AA	20030327	CA 2002-2465978	20020912
	EP 1429765	A2	20040623	EP 2002-763627	20020912
	R:				
	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK				
	BR 2002012510	A	20040824	BR 2002-12510	20020912
	JP 2005508905	T2	20050407	JP 2003-528544	20020912
	JP 2005255683	A2	20050922	JP 2005-80310	20050318
PRAI	US 2001-322402P	P	20010914		
	US 2002-391728P	P	20020626		
	JP 2003-528544	A3	20020912		
	WO 2002-US29017	W	20020912		
OS	MARPAT 138:271705				
GI					



AB The invention relates to triazines (shown as I; variables defined below; e.g. 4-[[[4-amino-6-(2-indanylamino)-[1,3,5]triazin-2-ylamino]methyl]-N-(2-aminophenyl)benzamide) and Cy3-X1-Ar2-(C(R5):C(R6))qC(O)NH-Ay2 (II; variables defined below; e.g.), many of which are N-(o-aminophenyl)carboxamides, as inhibitors of histone deacetylase (data included for many I and II). The invention provides compds. and methods for inhibiting histone deacetylase enzymic activity. The invention also provides compns. and methods for treating cell proliferative diseases and conditions. Antineoplastic effects of some I and II are illustrated for colorectal, pulmonary and pancreatic neoplasms; also the combined antineoplastic effect of histone deacetylase inhibitors and histone deacetylase antisense oligonucleotides on tumor cells in vivo was demonstrated. For I: R³ and R⁴ = H, L¹, Cy¹ and -L¹-Cy¹ (L¹ = C¹-C⁶ alkyl, C²-C⁶

heteroalkyl, or C3-C6 alkenyl; Cy1 = cycloalkyl, aryl, heteroaryl, or heterocyclyl) or R3 and R4 are taken together with the adjacent N atom to form a 5-, 6-, or 7-membered ring, wherein the ring atoms = C, O, S, and N, and wherein the ring is optionally substituted, and optionally forms part of a bicyclic ring system, or is optionally fused to one or two aryl or heteroaryl rings, or to one or two saturated or partially unsatd. cycloalkyl or heterocyclic rings, each of which rings and ring systems is optionally substituted. Y1 = -N(R1)(R2), -CH2-C(O)-N(R1)(R2), halogen, and H (R1 and R2 = H, L1, Cy1, and -L1-Cy1). Y2 = chemical bond or N(R0) (R0 = H, alkyl, aryl, aralkyl, and acyl); Ak1 = C1-C6 alkylene, C1-C6-heteroalkylene (preferably, in which one -CH2- is replaced with -NH-, and more preferably -NH-CH2), C2-C6 alkenylene or C2-C6 alkynylene; Ar1 = arylene or heteroarylene, either of which is optionally substituted; and Z1 = C(O)NH-Ay1 and CH:CHC(O)NH-Ay1 (Ay1 = aryl or heteroaryl, each of which is optionally substituted). For II: Cy2 = cycloalkyl, aryl, heteroaryl, or heterocyclyl; X1 = covalent bond, M1-L2-M1, and L2-M2-L2 (L2 = chemical bond, C1-C4 alkylene, C2-C4 alkenylene, and C2-C4 alkynylene, provided that L2 is not a chemical bond when X1 is M1-L2-M1; M1 = -O-, -N(R7)-, -S-, -S(O)-, S(O)2-, -S(O)2N(R7)-, -N(R7)S(O)2-, -C(O)-, -C(O)NH-, -NHC(O)-, -NHC(O)-O- and -OC(O)NH- (R7 = H, alkyl, aryl, aralkyl, acyl, heterocyclyl, and heteroaryl); and M2 = M1, heteroarylene, and heterocyclylene, either of which rings is optionally substituted). Ar2 = arylene or heteroarylene, each of which is optionally substituted; R5 and R6 = H, alkyl, aryl, and aralkyl; q is 0 or 1; and Ay2 is a 5-6 membered cycloalkyl, heterocyclyl, or heteroaryl substituted with an amino or hydroxy moiety (preferably these groups are ortho to the amide N to which Ay2 is attached) and further optionally substituted; provided that when Cy2 is naphthyl, X1 is -CH2-, Ar2 is Ph, R5 and R6 are H, and q is 0 or 1, Ay2 is not Ph or o-hydroxyphenyl. Although the methods of preparation are not claimed, hundreds of example prepns. are included.

L13 ANSWER 7 OF 8 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2002:907188 CAPLUS

DN 138:1673

TI **Inhibitors of histone deacetylase and their therapeutic use**

IN Curtin, Michael L.; Dai, Yujia; Davidsen, Steven K.; Frey, Robin R.; Guo, Yan; Heyman, Howard R.; Holms, James H.; Ji, Zhiqin; Michaelides, Michael R.; Vasudevan, Anil; Wada, Carol K.

PA USA

SO U.S. Pat. Appl. Publ., 49 pp.

CODEN: USXXCO

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2002177594	A1	20021128	US 2001-45747	20011026
PRAI	US 2001-275770P	P	20010314		
	US 2001-308435P	P	20010726		

OS MARPAT 138:1673

AB Compds. having the formula (R4L2)nL1CR1R2R3 (n = 1,2; L1 = alkenylene, alkylene, alkynylene, cycloalkylene, heteroalkylene, alkylene-CONR5-alkylene, alkylene-O-alkylene; L2 = bond, C2-alkenylene, O, S, SO2, OC(:O)NR5, NR6C:O, C(:O)NR6, SO2NR6, NR6SO2, C(:N)O, NR6C(:O)NR6, C(:O)NR6NR6C:O; R1 = alkanoyl, alkoxycarbonyl, aminocarbonyl, carboxy, haloalkyl, heterocycle; R2,R3 = OH or R2,R3 together = oxo; R4 = alkoxyalkyl, alkyl, aryl, arylalkyl, cycloalkyl, (cycloalkyl)alkyl, heterocycle, (heterocycle)alkyl; R5,R6 = hydrogen, alkyl, aryl, arylalkyl; R4,R6 and N to which they are attached = heterocycle) or therapeutically acceptable salts thereof, are histone **deacetylase (HDAC) inhibitors**. Preparation of the compds., compns. containing the compds., and treatment of diseases using the compds. are disclosed. Thus, more than 200 histone **deacetylase inhibitors** (no data) were

synthesized.

L13 ANSWER 8 OF 8 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2002:449627 CAPLUS

DN 137:33319

TI Preparation of N-aryl, N-arylalkyl, and N-heterocyclylnonanamide and -octanamide derivatives and related compounds as inhibitors of histone deacetylase

IN Curtin, Michael L.; Dai, Yujia; Davidsen, Steven K.; Frey, Robin R.; Guo, Yan; Heyman, Howard R.; Holms, James H.; Ji, Zhiqin; Michaelides, Michael R.; Vasudevan, Anil; Wada, Carol K.

PA Abbott Laboratories, USA

SO PCT Int. Appl., 111 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002046129	A2	20020613	WO 2001-US50931	20011026
	WO 2002046129	A3	20030116		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
	US 2002103192	A1	20020801	US 2001-808389	20010314
	AU 2002043402	A5	20020618	AU 2002-43402	20011026
PRAI	US 2000-697387	A	20001026		
	US 2001-808389	A	20010314		
	WO 2001-US50931	W	20011026		

OS MARPAT 137:33319

AB Compds. having the formula (R4-L2)nL1-CR1R2R3 or therapeutically acceptable salts thereof [wherein n = 1, 2; L1 = alkenylene, alkylene, alkynylene, cycloalkylene, heteroalkylene, (alkylene)-C(O)N(R5)-(alkylene), (alkylene)-O-(alkylene) (wherein each group is drawn with its left-hand end being the end which attaches to L2, and its right-hand end being the end which attaches to the carbon substituted with R1, R2, and R3); L2 =, C2 alkenylene, O, S, SO2, OC(O)NR5, N(R6)C(O), C(O)N(R6), SO2N(R6), N(R6)SO2, C:N-O, N(R6)C(O)N(R6), and C(O)N(R6)N(R6)C(O) (wherein each group is drawn with its left-hand end being the end which attaches to R4, and its right-hand end being the end which attaches to L1); R1 is selected from the group consisting of alkanoyl, alkoxycarbonyl, CONH2, CO2H, haloalkyl, heterocyclyl (wherein the heterocycle is selected from the group consisting of oxazolyl, dihydrooxazolyl, oxadiazolyl, and tetrazolyl); R2 = R3 = HO; or R2 and R3 together are oxo; R4 = alkoxyalkyl, alkyl, aryl, arylalkyl, cycloalkyl, cycloalkylalkyl, heterocycle, heterocyclylalkyl; R5, R6 = H, alkyl, aryl, arylalkyl; or R5 and R6, together with the nitrogen atom to which they are attached, form a heterocycle selected from the group consisting of (un)substituted morpholinyl, piperazinyl, piperidinyl, and thiomorpholinyl], which are histone deacetylase (HDAC) inhibitors (no data), are prepared These compds. are used for the treatment of diseases, possibly e.g. several human cancers associated with malfunction in histone deacetylases. Thus, a mixture of 9,9,9-trifluoro-8-oxononanoic acid (50 mg, 0.22 mmol), HOBt (30 mg, 0.22 mmol), carbodiimide PS resin (720 mg), and 4-phenyl-1,3-thiazol-2-amine (0.27 mmol) in DMF (5 mL) at room temperature was agitated in a Quest 210 parallel synthesizer for 18 h, treated with trisamine PS resin (220 mg), and agitated for 2 h. The solution was decanted, the resin was rinsed with dichloromethane, and the combined

solns. were concentrated, followed by purification using preparative HPLC with
a gradient system of 0 to 95 % over 10 min of MeCN (containing 0.1% CF₃CO₂H) in water to give 9,9,9-trifluoro-8-oxo-N-(4-phenyl-1,3-thiazol-2-yl)nonanamide.

=> analyze l13

ENTER ANSWER NUMBER OR RANGE (1-):1-8

ENTER DISPLAY CODE (TI) OR ?:rn

L14 ANALYZE L13 1-8 RN : 2801 TERMS

=> fil reg

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	192.65	366.68

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE ENTRY	TOTAL SESSION
CA SUBSCRIBER PRICE	-41.25	-41.25

FILE 'REGISTRY' ENTERED AT 16:18:26 ON 16 MAY 2006
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2006 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 15 MAY 2006 HIGHEST RN 884382-45-0
DICTIONARY FILE UPDATES: 15 MAY 2006 HIGHEST RN 884382-45-0

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH January 6, 2006

Please note that search-term pricing does apply when conducting SmartSELECT searches.

*
* The CA roles and document type information have been removed from *
* the IDE default display format and the ED field has been added, *
* effective March 20, 2005. A new display format, IDERL, is now *
* available and contains the CA role and document type information. *
*

Structure search iteration limits have been increased. See HELP SLIMITS for details.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/ONLINE/UG/regprops.html>

=> s l14

L15 2801 L14

=> d his

(FILE 'HOME' ENTERED AT 16:04:47 ON 16 MAY 2006)

FILE 'REGISTRY' ENTERED AT 16:05:20 ON 16 MAY 2006

L1 STRUC
L2 0 S L1
L3 2 S L1 FUL
L4 STRUC
L5 3 S L4
L6 3 S L5 NOT L3

FILE 'CAPLUS' ENTERED AT 16:10:12 ON 16 MAY 2006

L7 1 S L6
L8 2 S L3
L9 41 S (DEACETYLASE(L) INHIBITOR?) AND PIPERIDIN?
L10 41 S L9 NOT (L7 OR L8)
L11 23 S L10 AND (PYRROL? OR PYRAZO? OR OXAZO? OR FURAN? OR THIEN?)
L12 10 S L10 AND BENZIMIDAZ?
L13 8 S L12 AND L11
L14 ANALYZE L13 1-8 RN : 2801 TERMS


FILE 'REGISTRY' ENTERED AT 16:18:26 ON 16 MAY 2006

L15 2801 S L14

=> d 14

L4 HAS NO ANSWERS

L4 STR



VAR G1=13/14

NODE ATTRIBUTES:

DEFAULT MLEVEL IS ATOM

DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RSPEC 16 9

NUMBER OF NODES IS 28

STEREO ATTRIBUTES: NONE

=> search 14

ENTER TYPE OF SEARCH (SSS), CSS, FAMILY, OR EXACT:sss

ENTER SCOPE OF SEARCH (SAMPLE), FULL, RANGE, OR SUBSET:subset

ENTER SUBSET L# OR (END):l15

ENTER SUBSET SEARCH SCOPE - SAMPLE, FULL, RANGE, OR (END):ful

FULL SUBSET SEARCH INITIATED 16:18:58 FILE 'REGISTRY'

FULL SUBSET SCREEN SEARCH COMPLETED - 0 TO ITERATE

100.0% PROCESSED 0 ITERATIONS
SEARCH TIME: 00.00.01

0 ANSWERS

L16 0 SEA SUB=L15 SSS FUL L4

=> s l15 and benzimidazo?
329429 BENZIMIDAZO?


L17 81 L15 AND BENZIMIDAZO?

=> s l17 and piperidin?
946038 PIPERIDIN?

L18 2 L17 AND PIPERIDIN?

=> d 1-2


L18 ANSWER 1 OF 2 REGISTRY COPYRIGHT 2006 ACS on STN
RN 603986-54-5 REGISTRY
ED Entered STN: 14 Oct 2003
CN Benzamide, 2-[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]-N-hydroxy-3-nitro- (9CI) (CA INDEX NAME)
FS 3D CONCORD
MF C19 H19 N5 O5
CI COM
SR CA
LC STN Files: CA, CAPLUS, TOXCENTER



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

2 REFERENCES IN FILE CA (1907 TO DATE)
2 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L18 ANSWER 2 OF 2 REGISTRY COPYRIGHT 2006 ACS on STN
RN 603986-38-5 REGISTRY
ED Entered STN: 14 Oct 2003
CN Benzamide, 4-[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]-N-hydroxy-3-nitro- (9CI) (CA INDEX NAME)
FS 3D CONCORD
MF C19 H19 N5 O5
CI COM
SR CA
LC STN Files: CA, CAPLUS, TOXCENTER



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

2 REFERENCES IN FILE CA (1907 TO DATE)
2 REFERENCES IN FILE CAPLUS (1907 TO DATE)

=> fil reg

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	180.70	547.38
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE ENTRY	TOTAL SESSION
CA SUBSCRIBER PRICE	0.00	-41.25

FILE 'REGISTRY' ENTERED AT 16:19:45 ON 16 MAY 2006
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2006 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file
provided by InfoChem.

STRUCTURE FILE UPDATES: 15 MAY 2006 HIGHEST RN 884382-45-0
DICTIONARY FILE UPDATES: 15 MAY 2006 HIGHEST RN 884382-45-0

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH January 6, 2006

Please note that search-term pricing does apply when
conducting SmartSELECT searches.

*
* The CA roles and document type information have been removed from *
* the IDE default display format and the ED field has been added, *
* effective March 20, 2005. A new display format, IDERL, is now *
* available and contains the CA role and document type information. *
*

Structure search iteration limits have been increased. See HELP SLIMITS
for details.

REGISTRY includes numerically searchable data for experimental and
predicted properties as well as tags indicating availability of
experimental property data in the original document. For information
on property searching in REGISTRY, refer to:

<http://www.cas.org/ONLINE/UG/regprops.html>

=> s l18

946038 PIPERIDIN?

L19 2 L17 AND PIPERIDIN?

=> d bib abs hitstr

'BIB' IS NOT A VALID FORMAT FOR FILE 'REGISTRY'

'ABS' IS NOT A VALID FORMAT FOR FILE 'REGISTRY'

'HITSTR' IS NOT A VALID FORMAT FOR FILE 'REGISTRY'

The following are valid formats:

Substance information can be displayed by requesting individual fields or predefined formats. The predefined substance formats are: (RN = CAS Registry Number)

REG - RN
SAM - Index Name, MF, and structure - no RN
FIDE - All substance data, except sequence data
IDE - FIDE, but only 50 names
SQIDE - IDE, plus sequence data
SQIDE3 - Same as SQIDE, but 3-letter amino acid codes are used
SQD - Protein sequence data, includes RN
SQD3 - Same as SQD, but 3-letter amino acid codes are used
SQN - Protein sequence name information, includes RN

CALC - Table of calculated properties
EPROP - Table of experimental properties
PROP - EPROP and CALC

Any CA File format may be combined with any substance format to obtain CA references citing the substance. The substance formats must be cited first. The CA File predefined formats are:

ABS -- Abstract
APPS -- Application and Priority Information
BIB -- CA Accession Number, plus Bibliographic Data
CAN -- CA Accession Number
CBIB -- CA Accession Number, plus Bibliographic Data (compressed)
IND -- Index Data
IPC -- International Patent Classification
PATS -- PI, SO
STD -- BIB, IPC, and NCL

IABS -- ABS, indented, with text labels
IBIB -- BIB, indented, with text labels
ISTD -- STD format, indented

OBIB ----- AN, plus Bibliographic Data (original)
OIBIB ----- OBIB, indented with text labels

SBIB ----- BIB, no citations
SIBIB ----- IBIB, no citations

The ALL format gives FIDE BIB ABS IND RE, plus sequence data when it is available.

The MAX format is the same as ALL.

The IALL format is the same as ALL with BIB ABS and IND indented, with text labels.

For additional information, please consult the following help messages:

HELP DFIELDS -- To see a complete list of individual display fields.
HELP FORMATS -- To see detailed descriptions of the predefined formats.
ENTER DISPLAY FORMAT (IDE):bib abs hitstr
'BIB' IS NOT A VALID FORMAT FOR FILE 'REGISTRY'
'ABS' IS NOT A VALID FORMAT FOR FILE 'REGISTRY'
'HITSTR' IS NOT A VALID FORMAT FOR FILE 'REGISTRY'

The following are valid formats:

Substance information can be displayed by requesting individual fields or predefined formats. The predefined substance formats are: (RN = CAS Registry Number)

REG - RN
 SAM - Index Name, MF, and structure - no RN
 FIDE - All substance data, except sequence data
 IDE - FIDE, but only 50 names
 SQIDE - IDE, plus sequence data
 SQIDE3 - Same as SQIDE, but 3-letter amino acid codes are used
 SQD - Protein sequence data, includes RN
 SQD3 - Same as SQD, but 3-letter amino acid codes are used
 SQN - Protein sequence name information, includes RN

CALC - Table of calculated properties
 EPROP - Table of experimental properties
 PROP - EPROP and CALC

Any CA File format may be combined with any substance format to obtain CA references citing the substance. The substance formats must be cited first. The CA File predefined formats are:

ABS -- Abstract
 APPS -- Application and Priority Information
 BIB -- CA Accession Number, plus Bibliographic Data
 CAN -- CA Accession Number
 CBIB -- CA Accession Number, plus Bibliographic Data (compressed)
 IND -- Index Data
 IPC -- International Patent Classification
 PATS -- PI, SO
 STD -- BIB, IPC, and NCL

IABS -- ABS, indented, with text labels
 IBIB -- BIB, indented, with text labels
 ISTD -- STD format, indented

OBIB ----- AN, plus Bibliographic Data (original)
 OIBIB ----- OBIB, indented with text labels

SBIB ----- BIB, no citations
 SIBIB ----- IBIB, no citations

The ALL format gives FIDE BIB ABS IND RE, plus sequence data when it is available.

The MAX format is the same as ALL.

The IALL format is the same as ALL with BIB ABS and IND indented, with text labels.

For additional information, please consult the following help messages:

HELP DFIELDS -- To see a complete list of individual display fields.
 HELP FORMATS -- To see detailed descriptions of the predefined formats.
 ENTER DISPLAY FORMAT (IDE):end

=> d his

(FILE 'HOME' ENTERED AT 16:04:47 ON 16 MAY 2006)

FILE 'REGISTRY' ENTERED AT 16:05:20 ON 16 MAY 2006

L1 STRUC
 L2 0 S L1
 L3 2 S L1 FUL
 L4 STRUC
 L5 3 S L4
 L6 3 S L5 NOT L3

FILE 'CAPLUS' ENTERED AT 16:10:12 ON 16 MAY 2006

L7 1 S L6
L8 2 S L3
L9 41 S (DEACETYLASE(L) INHIBITOR?) AND PIPERIDIN?
L10 41 S L9 NOT (L7 OR L8)
L11 23 S L10 AND (PYRROL? OR PYRAZO? OR OXAZO? OR FURAN? OR THIEN?)
L12 10 S L10 AND BENZIMIDAZ?
L13 8 S L12 AND L11
L14 ANALYZE L13 1-8 RN : 2801 TERMS

FILE 'REGISTRY' ENTERED AT 16:18:26 ON 16 MAY 2006

L15 2801 S L14
L16 0 SEARCH L4 SSS SUB=L15 FUL
L17 81 S L15 AND BENZIMIDAZO?
L18 2 S L17 AND PIPERIDIN?

FILE 'REGISTRY' ENTERED AT 16:19:45 ON 16 MAY 2006

L19 2 S L18

=> s 118

L20 2 L18

=> d bib abs hitstr 1-2

L20 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2005:300395 CAPLUS

DN 142:355054

TI Preparation of amide derivatives as inhibitors of histone deacetylase

IN Moradei, Oscar; Paquin, Isabelle; Leit, Silvana; Frechette, Sylvie;
Vaisburg, Arkadii; Besterman, Jeffrey M.; Tessier, Pierre; Mallais, Tammy
C.

PA Methylgene, Inc., Can.

SO PCT Int. Appl., 559 pp.



CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	WO 2005030705	A1	20050407	WO 2004-US31591	20040924
	WO 2005030705	C2	20060420		
	W:				
	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH,				
	CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD,				
	GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,				
	LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI,				
	NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY,				
	TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW:				
	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM,				
	AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK,				
	EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE,				
	SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE,				
	SN, TD, TG				
PRAI	US 2003-505884P	P	20030924		
	US 2003-532973P	P	20031229		
	US 2004-561082P	P	20040409		
OS	MARPAT 142:355054				
GI					



AB Title compds. I [Ar1 = (un)saturated-, (un)substituted-mono or fused poly-cyclic hydrocarbyl optionally containing 1-4 heteroatoms per ring; R1 = (un)substituted-mono-, -bi-, -tri-cyclic-aryl or -heteroaryl; R2, R3, and R4 independently = H, halo, amino, etc.; R5 and R6 independently = H, alkyl, aryl, etc.; x = 0-1; Y = any pharmaceutically acceptable chemical moiety consisting of 1 to 50 atoms with provisions] and their pharmaceutically acceptable salts, are prepared and disclosed as inhibitors of histone deacetylase. Thus, e.g., II was prepared by Suzuki coupling of 2-bromo-2-nitro-phenylamine (preparation given) with 2-thiopheneboronic acid followed by carbonylation with 4-[3,4-dimethoxy-(phenylamino)-methyl]benzoic acid (preparation given) and subsequent reduction. The inhibitory

capability of I towards antiproliferative activity of histone deacetylase enzyme was evaluated using 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide (MTT) assay and it revealed that certain compds. of the invention had MTT IC 50 values in the range of below 1 up to 20 μ M. I as histone deacetylase inhibitors should prove useful in the treatment of diseases such as, but not limited to, cell proliferative disease, protozoal disease, and fungal disease.


IT 603986-38-5P 603986-54-5P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of amide derivs. as inhibitors of histone deacetylase)


RN 603986-38-5 CAPLUS

CN Benzamide, 4-[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]-N-hydroxy-3-nitro- (9CI) (CA INDEX NAME)



RN 603986-54-5 CAPLUS

CN Benzamide, 2-[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]-N-hydroxy-3-nitro- (9CI) (CA INDEX NAME)



RE.CNT 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L20 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2006 ACS on STM

AN 2005:300394 CAPLUS

DN 142:373563

TI Preparation of amide derivatives as inhibitors of histone deacetylase

IN Moradei, Oscar; Paquin, Isabelle; Leit, Silvana; Frechette, Sylvie; Vaisburg, Arkadii; Besterman, Jeffrey M.; Tessier, Pierre; Mallais, Tammy C.

PA Methylgene, Inc., Can.

SO PCT Int. Appl., 389 pp.



CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2005030704	A1	20050407	WO 2004-US31590	20040924
	W:				
	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW:				
	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
PRAI	US 2003-505884P	P	20030924		
	US 2003-532973P	P	20031229		
	US 2004-561082P	P	20040409		
OS	MARPAT 142:373563				
GI					



AB Title compds. I [Ar1 = (un)saturated-, (un)substituted-mono or fused poly-cyclic hydrocarbonyl optionally containing 1-4 heteroatoms per ring; R1 = (un)substituted-mono-, -bi-, -tri-cyclic-aryl or -heteroaryl; R2, R3, and R4 independently = H, halo, amino, etc.; R5 and R6 independently = H, alkyl, aryl, etc.; x = 0-1; Y = any pharmaceutically acceptable chemical moiety consisting of 1 to 50 atoms with provisions] and their pharmaceutically acceptable salts, are prepared and disclosed as inhibitors of histone deacetylase. Thus, e.g., II was prepared by Suzuki coupling of 2-bromo-2-nitro-phenylamine (preparation given) with 2-thiopheneboronic acid followed by carbonylation with 4-[3,4-dimethoxy-(phenylamino)-methyl]benzoic acid (preparation given) and subsequent reduction. The inhibitory

capability of I towards antiproliferative activity of histone deacetylase enzyme was evaluated using 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide (MTT) assay and it revealed that certain compds. of the invention had MTT IC 50 values in the range of below 1 up to 20 μ M. I as histone deacetylase inhibitors should prove useful in the treatment of diseases such as, but not limited to, cell proliferative disease, protozoal disease, and fungal disease.


IT 603986-38-5P 603986-54-5P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of amide derivs. as inhibitors of histone deacetylase)


RN 603986-38-5 CAPLUS

CN Benzamide, 4-[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]-N-hydroxy-3-nitro- (9CI) (CA INDEX NAME)



RN 603986-54-5 CAPLUS


CN Benzamide, 2-[4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-1-piperidinyl]-N-hydroxy-3-nitro- (9CI) (CA INDEX NAME)



RE.CNT 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

AN 2004:799454 CAPLUS
 DN 141:291229
 TI Histone deacetylase inhibitors
 IN Bressi, Jerome C.; Brown, Jason W.; Cao, Sheldon X.; Gangloff, Anthony R.;
 Jennings, Andrew J.; Stafford, Jeffrey A.; Vu, Phong H.; Xiao, Xiao-Yi
 PA Syrrx, Inc., USA
 SO PCT Int. Appl., 276 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004082638	A2	20040930	WO 2004-US8342	20040317
	WO 2004082638	A3	20050506		
	W:				
	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW:				
	BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	CA 2518318	AA	20040930	CA 2004-2518318	20040317
	US 2004254220	A1	20041216	US 2004-803575	20040317
	US 2004266769	A1	20041230	US 2004-803344	20040317
	US 2005137232	A1	20050623	US 2004-803580	20040317
	EP 1608628	A2	20051228	EP 2004-757631	20040317
	R:				
	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK				
PRAI	US 2003-455437P	P	20030317		
	US 2003-531203P	P	20031219		
	WO 2004-US8342	W	20040317		
OS	MARPAT 141:291229				
AB	Compds. that may be used to inhibit histone deacetylase are disclosed. Thus, 119 compds. were prepared which exhibited better than 1000 nM IC50 against HDAC1, HDAC2, HDAC6, and HDAC8 (suberanilohydroxamic acid showed an IC50 of 63 nM in this assay). Many of these compds. were 3-[3-(1-substituted-1H-benzimidazol-2-yl)phenyl]acrylic acids and N-hydroxy- [3-(1-substituted-1H-benzimidazol-2-yl)phenyl]acrylamides.				
IT	758693-30-0 758694-08-5 758694-10-9				
	RL: BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)				
	(histone deacetylase inhibitors)				
RN	758693-30-0 CAPLUS				
CN	2-Propenamide, 3-[3-[1-(1-ethyl-3-piperidinyl)-1H-benzimidazol-2-yl]phenyl]-N-hydroxy- (9CI) (CA INDEX NAME)				




RN 758694-08-5 CAPLUS

CN 2-Propenamide, 3-[3-[1-[(3R)-1-ethyl-3-piperidinyl]-5,6-difluoro-1H-benzimidazol-2-yl]phenyl]-N-hydroxy- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Double bond geometry unknown.




RN 758694-10-9 CAPLUS

CN 2-Propenamide, 3-[3-[1-[(3R)-1-ethyl-3-piperidinyl]-5,6,7-trifluoro-1H-benzimidazol-2-yl]phenyl]-N-hydroxy- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

Double bond geometry unknown.



AN 2002:220554 CAPLUS
 DN 136:262995
 TI Preparation of hydroxamic acids as deacetylase inhibitors
 IN Bair, Kenneth Walter; Green, Michael A.; Perez, Lawrence B.; Remiszewski, Stacy W.; Sambucetti, Lidia; Versace, Richard William; Sharma, Sushil Kumar
 PA Novartis AG, Switz.; Novartis-Erfindungen Verwaltungsgesellschaft mbH; Novartis Pharma GmbH
 SO PCT Int. Appl., 96 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002022577	A2	20020321	WO 2001-EP10037	20010830
	WO 2002022577	A3	20020906		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
	CA 2420899	AA	20020321	CA 2001-2420899	20010830
	AU 2001082129	A5	20020326	AU 2001-82129	20010830
	BR 2001013669	A	20030603	BR 2001-13669	20010830
	EP 1318980	A2	20030618	EP 2001-960717	20010830
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
	JP 2004509105	T2	20040325	JP 2002-526830	20010830
	NZ 524365	A	20041126	NZ 2001-524365	20010830
	US 2003018062	A1	20030123	US 2001-944275	20010831
	US 6552065	B2	20030422		
	US 2004024067	A1	20040205	US 2002-299518	20021116
	ZA 2003001423	A	20040421	ZA 2003-1423	20030221
	NO 2003000867	A	20030225	NO 2003-867	20030225
	US 2005085507	A1	20050421	US 2004-984501	20041109
PRAI	US 2000-229943P	P	20000901		
	US 2001-292232P	P	20010518		
	US 2001-307490P	P	20010724		
	WO 2001-EP10037	W	20010830		
	US 2001-944275	A1	20010831		
	US 2002-299518	A1	20021116		
OS	MARPAT 136:262995				
GI					



II

AB The title compds. [I; R1 = H, halo, alkyl; R2 = H, alkyl, cycloalkyl, etc.; R3, R4 = H, alkyl, acyl, acylamino; or R3 and R4 together with the carbon atom to which they are bound = CO, CS, C:NR8; or R2 together with the N atom to which is bound and R3 together with the C atom to which it is bound form heterocycloalkyl, heteroaryl, etc.; R5 = H, alkyl, aryl, etc.; n1-n3 = 0-6; X, Y = H, halo, alkyl, etc.; R8 = H, alkyl, aryl, etc.] which are deacetylase inhibitors and therefore suitable for pharmaceutical compns. having anti-proliferative properties, were prepared E.g., a 3-step synthesis of II, starting with 4-formylcinnamic acid, was given. The exemplified compds. I showed IC50 of 0.005-0.5 μ M against HDA.

IT 404949-04-8P


RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of hydroxamic acids as deacetylase inhibitors)

RN 404949-04-8 CAPLUS

CN 1-Piperidinecarboxylic acid, 4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-, 2-[[[4-[(1E)-3-(hydroxyamino)-3-oxo-1-propenyl]phenyl]methyl][2-(1H-indol-3-yl)ethyl]amino]ethyl ester (9CI) (CA INDEX NAME)


Double bond geometry as shown.



=> d l1

L1 HAS NO ANSWERS

L1 STR



VAR G1=O/S/C/N

VAR G2=C/N

REP G3=(0-10) CH

VAR G4=O/CY

NODE ATTRIBUTES:

DEFAULT MLEVEL IS ATOM

DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RSPEC 6 4

NUMBER OF NODES IS 16

STEREO ATTRIBUTES: NONE

=> s l1 ful

FULL SEARCH INITIATED 17:07:09 FILE 'REGISTRY'

FULL SCREEN SEARCH COMPLETED - 731027 TO ITERATE

98.2% PROCESSED 718034 ITERATIONS

2 ANSWERS

100.0% PROCESSED 731027 ITERATIONS

2 ANSWERS

SEARCH TIME: 00.00.29

L3 2 SEA SSS FUL L1

=> d 1-2

L3 ANSWER 1 OF 2 REGISTRY COPYRIGHT 2006 ACS on STN

RN 709654-55-7 REGISTRY

ED Entered STN: 14 Jul 2004


CN 1H-Pyrazole-5-carboxamide, N-[2'-[[[(1,1-dimethylethyl)amino]sulfonyl][1,1'-biphenyl]-4-yl]-3-methyl-1-(3-piperidinyl)-(9CI) (CA INDEX NAME)

FS 3D CONCORD

MF C26 H33 N5 O3 S

SR CA


LC STN Files: CA, CAPLUS, CASREACT



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

1 REFERENCES IN FILE CA (1907 TO DATE)
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

L3 ANSWER 2 OF 2 REGISTRY COPYRIGHT 2006 ACS on STN
RN 629610-26-0 REGISTRY
ED Entered STN: 22 Dec 2003
CN 1H-Pyrazole-5-carboxamide, N-[2'-(aminosulfonyl)[1,1'-biphenyl]-4-yl]-3-methyl-1-(3-piperidinyl)- (9CI) (CA INDEX NAME)
FS 3D CONCORD
MF C22 H25 N5 O3 S
SR CA
LC STN Files: CA, CAPLUS, CASREACT



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

2 REFERENCES IN FILE CA (1907 TO DATE)
2 REFERENCES IN FILE CAPLUS (1907 TO DATE)

=> fil caplus
COST IN U.S. DOLLARS
FULL ESTIMATED COST

SINCE FILE	TOTAL
ENTRY	SESSION
174.26	174.47

FILE 'CAPLUS' ENTERED AT 17:08:14 ON 17 MAY 2006
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the

American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 17 May 2006 VOL 144 ISS 21
FILE LAST UPDATED: 16 May 2006 (20060516/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:


<http://www.cas.org/infopolicy.html>

=> s 13

L4 2 L3


=> d bib abs hitstr 1-2

L4 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2006 ACS on STN
AN 2006:165187 CAPLUS
DN 144:304521
TI Comparative study of factor Xa inhibitors using molecular docking/SVM/HQSAR/3D-QSAR methods
AU Sun, Jing; Chen, Hai Feng; Xia, Hai Rong; Yao, Jian Hua; Fan, Bo Tao
CS Laboratory of Computer Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, Shanghai, 200032, Peop. Rep. China
SO QSAR & Combinatorial Science (2006), 25(1), 25-45
CODEN: QCSSAU; ISSN: 1611-020X
PB Wiley-VCH Verlag GmbH & Co. KGaA
DT Journal
LA English
AB The binding modes of a group of Factor Xa (fXa) inhibitors were studied using FlexX. CoMFA, CoMSIA, HQSAR and SVM models for inhibition potency were constructed with the conformers obtained from the mol. docking. 3D-QSAR models for oral bioavailability were also constructed with the subset inhibitors. The results show that these models possess good prediction ability. The influence of substituents for the activity and oral bioavailability were explored by comparing the constructed 3D-QSAR models. We found that some substituents have consistent effects on inhibition potency and oral bioavailability, but some have inconsistent effects. We observed equally that the different methods involved in this study, such as mol. docking, SVM, HQSAR and 3D-QSAR models, could be used not only for the prediction, but they are also complementary each to other. They are helpful for better understanding the interaction mechanism between inhibitors and fXa receptor.
IT 629610-26-0
RL: PAC (Pharmacological activity); PKT (Pharmacokinetics); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (comparative study of factor Xa inhibitors using mol. docking/SVM/HQSAR/QSAR methods)
RN 629610-26-0 CAPLUS
CN 1H-Pyrazole-5-carboxamide, N-[2'-(aminosulfonyl)[1,1'-biphenyl]-4-yl]-3-methyl-1-(3-piperidinyl)- (9CI) (CA INDEX NAME)



RE.CNT 46 THERE ARE 46 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2006 ACS on STN
AN 2003:784065 CAPLUS
DN 140:12453
TI Structure-based design of novel guanidine/benzamidine mimics: potent and orally bioavailable factor Xa inhibitors as novel anticoagulants
AU Lam, Patrick Y. S.; Clark, Charles G.; Li, Renhua; Pinto, Donald J. P.; Orwat, Michael J.; Galembo, Robert A.; Fevig, John M.; Teleha, Christopher A.; Alexander, Richard S.; Smallwood, Angela M.; Rossi, Karen A.; Wright, Matthew R.; Bai, Stephen A.; He, Kan; Luetzgen, Joseph M.; Wong, Pancras C.; Knabb, Robert M.; Wexler, Ruth R.
CS Bristol-Myers Squibb Company, Princeton, NJ, 08542-5400, USA
SO Journal of Medicinal Chemistry (2003), 46(21), 4405-4418
CODEN: JMCMAR; ISSN: 0022-2623
PB American Chemical Society
DT Journal
LA English
OS CASREACT 140:12453
AB As part of an ongoing effort to prepare orally active factor Xa inhibitors using structure-based drug design techniques and mol. recognition principles, a systematic study has been performed on the pharmacokinetic profile resulting from replacing the benzamidine in the P1 position with less basic benzamidine mimics or neutral residues. It is demonstrated that lowering the pKa of the P1 ligand resulted in compds. (3-benzylamine, 15a; 1-aminoisoquinoline, 24a; 3-aminobenzisoxazole, 23a; 3-phenylcarboxamide, 22b; and 4-methoxyphenyl, 22a) with improved pharmacokinetic features mainly as a result of decreased clearance, increased volume of distribution, and enhanced oral absorption. This work resulted in a series of potent and orally bioavailable factor Xa inhibitors that ultimately led to the discovery of SQ311, 24a. SQ311, which utilizes a 1-aminoisoquinoline as the P1 ligand, inhibits factor Xa with a Ki of 0.33 nM and demonstrates both good in vivo antithrombotic efficacy and oral bioavailability.
IT 629610-26-0P
RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(guanidine/benzamidine mimics as potent and orally bioavailable factor Xa inhibitors and anticoagulants)
RN 629610-26-0 CAPLUS
CN 1H-Pyrazole-5-carboxamide, N-[2'-(aminosulfonyl)[1,1'-biphenyl]-4-yl]-3-methyl-1-(3-piperidinyl)- (9CI) (CA INDEX NAME)




IT 709654-55-7

RL: RCT (Reactant); SPN (Synthetic preparation)

(guanidine/benzamidine mimics as potent and orally bioavailable factor
Xa inhibitors and anticoagulants)


RN 709654-55-7 CAPLUS

CN 1H-Pyrazole-5-carboxamide, N-[2'-[[[(1,1-dimethylethyl)amino]sulfonyl][1,1'-
biphenyl]-4-yl]-3-methyl-1-(3-piperidinyl)- (9CI) (CA INDEX NAME)



RE.CNT 44 THERE ARE 44 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

> d l27
 L27 HAS NO ANSWERS
 L27 STR



REP G1=(0-10) CH
 VAR G2=O/S/N
 VAR G3=12/13/14
 NODE ATTRIBUTES:
 DEFAULT MLEVEL IS ATOM
 DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:
 RSPEC 8
 NUMBER OF NODES IS 21

STEREO ATTRIBUTES: NONE

=> s l27 ful
 FULL SEARCH INITIATED 16:29:13 FILE 'REGISTRY'
 FULL SCREEN SEARCH COMPLETED - 681 TO ITERATE

100.0% PROCESSED 681 ITERATIONS 7 ANSWERS
 SEARCH TIME: 00.00.01


L29 7 SEA SSS FUL L27

=> d scan


L29 7 ANSWERS REGISTRY COPYRIGHT 2006 ACS on STN
 IN 1H-Benzimidazole-2-butanamide, α -amino-N-hydroxy-1-(1-methyl-4-piperidinyl)-, (α S)-, trifluoroacetate (salt) (9CI)
 MF C17 H25 N5 O2 . x C2 H F3 O2

CM 1

Absolute stereochemistry.



CM 2



HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):0

=> fil caplus

COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
170.90	736.46

FULL ESTIMATED COST

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE	TOTAL
ENTRY	SESSION
0.00	-42.75

CA SUBSCRIBER PRICE

FILE 'CAPLUS' ENTERED AT 16:29:34 ON 16 MAY 2006

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 16 May 2006 VOL 144 ISS 21

FILE LAST UPDATED: 15 May 2006 (20060515/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/infopolicy.html>



=> s 129

L30 6 L29

=> d bib abs hitstr 1-6

L30 ANSWER 1 OF 6 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 2005:729627 CAPLUS
 DN 143:212171
 TI Preparation of hydroxamic acid derivatives as AGE generation inhibitors
 IN Kakuchi, Junji; Yamazaki, Toru; Obara, Kazumi; Yamato, Hideyuki
 PA Kureha Chemical Industry Company, Limited, Japan
 SO PCT Int. Appl., 215 pp.
 CODEN: PIXXD2
 DT Patent
 LA Japanese
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2005073180	A1	20050811	WO 2004-JP19512	20041227
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
PRAI	JP 2003-428901	A	20031225		
OS	MARPAT 143:212171				
GI					



AB Title compds. I [R1 = H, alkyl, etc.; A1, A2 = single bond, etc.; Q1 = -Y1-A3-R2, etc.; Y1 = O, etc.; A3 = single bond, etc.; R2 = alkyl, etc.] were prepared For example, reductive amination of EDCI mediated resin bound N α -BOC-ornithine hydroxamic acid with propionaldehyde using sodium cyanoborohydride followed by treatment with trifluoroacetic acid afforded compound II [X = dipropylamino] trifluoroacetic acid salt. In Maillard reaction inhibition assays, compound II [X = bis(4-methylbenzyl)amino] trifluoroacetic acid salt showed the activity of 100% at 0.1 mM. Compds. I are claimed useful as AGE generation inhibitors.

IT 862400-22-4P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES

(Uses)

(preparation of hydroxamic acid derivs. as AGE generation inhibitors)

RN 862400-22-4 CAPLUS


CN 1H-Benzimidazole-2-butanamide, α -amino-N-hydroxy-1-(1-methyl-4-piperidiny)-, (α S)-, trifluoroacetate (salt) (9CI) (CA INDEX NAME)

CM 1

CRN 862400-21-3

CMF C17 H25 N5 O2


Absolute stereochemistry.



CM 2

CRN 76-05-1

CMF C2 H F3 O2



RE.CNT 37 THERE ARE 37 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L30 ANSWER 2 OF 6 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2002:10123 CAPLUS

DN 136:64091

TI Method and system for predicting pharmacokinetic properties

IN Hattori, Kazunari; Shimada, Kaore; Uchiyama, Mamoru

PA Pfizer Inc., USA

SO Eur. Pat. Appl., 27 pp.

CODEN: EPXXDW

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 1167969	A2	20020102	EP 2001-304648	20010525
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
	US 2003069698	A1	20030410	US 2001-876767	20010607
	JP 2003014728	A2	20030115	JP 2001-179774	20010614
PRAI	US 2000-211864P	P	20000614		

AB This invention provides a method for predicting pharmacokinetic properties of mols. comprising the steps of: (a) preparing 2D-structures of mols. used as a training set; (b) constructing a 2D-fingerprint by counting the number


of structural descriptors that potentially relate to a pharmacokinetic property, either manually or automatically using internally developed macro; wherein said structural descriptors consist of predefined 20 to 80 atoms/fragments or substructures; (c) analyzing the obtained 2D-fingerprint by a statistical anal. method to correlate with the pharmacokinetic property of the mol. to yield a quant. structure-property relation (QSPR) model; and (d) calculating the pharmacokinetic property of a trial mol. using the above obtained QSPR model. A system for this invention is also provided. According to this method and system, it is possible to predict pharmacokinetic properties of mols. prior to synthesis, without labor-intensive and time-consuming experimentation.

IT 258286-85-0

RL: PKT (Pharmacokinetics); PRP (Properties); BIOL (Biological study)
(method and system for predicting pharmacokinetic properties)

RN 258286-85-0 CAPLUS

CN 1-Propanone, 3-amino-1-[1-[1-(1-phenylcycloheptyl)-4-piperidiny]-1H-benzimidazol-2-yl]- (9CI) (CA INDEX NAME)



L30 ANSWER 3 OF 6 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2001:573269 CAPLUS

DN 135:152805

TI Preparation of benzimidazoles as ORL1-receptor agonists for analgesics

IN Ito, Fumitaka; Noguchi, Hirohide; Ohashi, Yoriko; Shimokawa, Hirohisa

PA Pfizer Pharmaceutical Co., Ltd., Japan

SO Jpn. Kokai Tokkyo Koho, 39 pp.

CODEN: JKXXAF

DT Patent


LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	JP 2001213878	A2	20010807	JP 2000-396414	20001227
	JP 3392402	B2	20030331		
	EP 1122257	A1	20010808	EP 2000-311316	20001218
	EP 1122257	B1	20051012		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
	AT 306488	E	20051015	AT 2000-311316	20001218
	ES 2249237	T3	20060401	ES 2000-311316	20001218
	CA 2330092	AA	20010705	CA 2001-2330092	20010103
	CA 2330092	C	20050322		
	US 2002049212	A1	20020425	US 2001-753954	20010103
	US 6861425	B2	20050301		
	BR 2001000014	A	20010828	BR 2001-14	20010104
PRAI	US 2000-174542P	P	20000105		

OS MARPAT 135:152805

GI




AB Title compds. I [R1 = C3-11 cycloalkyl, C6-16 bicycloalkyl, C6-16 tricycloalkyl, C8-16 tetracycloalkyl, etc.; A = (un)substituted C1-7 alkyl, C2-5 alkenyl, C2-5 alkynyl, aryl, etc.; M = single bond, CH2, O, S, SO, SO2, CO, NH, etc.; Y = 4- to 12-membered bicyclic carbon ring, 4- to 12- membered bicyclic hetero ring, 5- to 17-membered spiro carbon ring, 5- to 17-membered spiro hetero ring; Z1-Z4 = (un)substituted C1-4 alkyl, C1-4 alkoxy, C1-4 alkylsulfonyl, C1-4 alkylcarbonyl, carboxy, etc.] or their salts are prepared Tert-Bu 3-[1-[1-(1-phenylcycloheptyl)-4-piperidinyl]-1H-benzimidazol-2-yl]-3,8-diazabicyclo[3.2.1]octane-8-carboxylate was treated with F3CCO2H in CH2Cl2 at room temperature for 0.5 h to give 77.6% 2-(3,8-diazabicyclo[3.2.1]oct-3-yl)-1-[1-(1-phenylcycloheptyl)-4-piperidinyl]-1H-benzimidazole HCl salt.

IT 352541-85-6P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (preparation of benzimidazoles as ORL1-receptor agonists for analgesics)

RN 352541-85-6 CAPLUS

CN 1H-Benzimidazole-2-carboxylic acid, 1-[1-(1-phenylcycloheptyl)-4-piperidinyl]-, methyl ester (9CI) (CA INDEX NAME)



L30 ANSWER 4 OF 6 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2000:117042 CAPLUS

DN 132:151821

TI Preparation of 2-substituted-1-piperidylbenzimidazoles as ORL1 receptor agonists.

IN Ito, Fumitaka; Noguchi, Hirohide; Kondo, Hiroshi

PA Pfizer Pharmaceuticals Inc., Japan; Pfizer Inc.

SO PCT Int. Appl., 127 pp.
 CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2000008013	A2	20000217	WO 1999-IB1239	19990705
	WO 2000008013	A3	20000323		

W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ,

TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ,
MD, RU, TJ, TM
RW: GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG

TW 513424	B	20021211	TW 1999-88110899	19990628
CA 2339621	AA	20000217	CA 1999-2339621	19990705
CA 2339621	C	20050405		
AU 9943859	A1	20000228	AU 1999-43859	19990705
AU 749166	B2	20020620		
EP 1102762	A2	20010530	EP 1999-926688	19990705
EP 1102762	B1	20021113		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
TR 200100403	T2	20010723	TR 2001-200100403	19990705
BR 9912778	A	20010925	BR 1999-12778	19990705
EE 200100075	A	20020617	EE 2001-75	19990705
JP 2002522431	T2	20020723	JP 2000-563646	19990705
JP 3367945	B2	20030120		
AT 227716	E	20021115	AT 1999-926688	19990705
PT 1102762	T	20030228	PT 1999-926688	19990705
ES 2185357	T3	20030416	ES 1999-926688	19990705
NZ 509299	A	20030530	NZ 1999-509299	19990705
US 6172067	B1	20010109	US 1999-369208	19990805
ZA 2001000900	A	20020603	ZA 2001-900	20010201
HR 2001000089	A1	20020228	HR 2001-89	20010202
HR 20010089	B1	20030430		
NO 2001000603	A	20010405	NO 2001-603	20010205
BG 105301	A	20011231	BG 2001-105301	20010301
US 2003109549	A1	20030612	US 2002-283604	20021030
PRAI WO 1998-IB1206	W	19980806		
WO 1999-IB1239	W	19990705		
US 1999-369208	A3	19990805		
US 2000-676245	B1	20000929		

OS MARPAT 132:151821
GI




AB Title compds. [I; R = (substituted) mono-, di-, tri-, or tetracycloalkyl; A = alkyl, haloalkyl, alkenyl, alkynyl, (substituted) phenylalkyl, aryl, heteroaryl, heterocyclyl; Y = H, halo, amino, SH, (substituted) alkyl-M, cycloalkyl-M, alkenyl-M, alkyl-NH-alkyl-M, dialkyl-N-alkyl-M, aryl-M, heterocyclyl-M, arylalkyl-M, etc.; M = bond, O, S, NH S, SO, SO₂, etc.; Z1-Z4 = H, halo, alkyl, haloalkyl, alkoxy, alkylsulfonyl, alkylcarbonyl, CO₂H, amino, H₂NCO, Ph, naphthyl, etc.], were prepared as ORL1 receptor agonists (no data). Thus, 2-chloro-1-[1-(1-phenylcycloheptyl)-4-piperidinyl]benzimidazole (preparation given) was stirred with MeNH₂ in MeOH in an autoclave at 110° for 6 h to give N-methyl-1-[1-(1-phenylcycloheptyl)-4-piperidinyl]-1H-benzimidazol-2-amine.

IT 258286-85-OP 258287-70-6P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(preparation of 2-substituted-1-piperidylbenzimidazoles as ORL1 receptor agonists)


RN 258286-85-0 CAPLUS

CN 1-Propanone, 3-amino-1-[1-[1-(1-phenylcycloheptyl)-4-piperidiny]-1H-benzimidazol-2-yl]- (9CI) (CA INDEX NAME)



RN 258287-70-6 CAPLUS

CN 1-Propanone, 3-amino-1-[1-[1-(1-phenylcycloheptyl)-4-piperidiny]-1H-benzimidazol-2-yl]-, dihydrochloride (9CI) (CA INDEX NAME)



●2 HCl

L30 ANSWER 5 OF 6 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1997:51439 CAPLUS

DN 126:89269

TI Preparation of heterocyclic compounds as cholesterol acyltransferase inhibitors

IN Natsukari, Hideaki; Ishimaru, Takenori; Doi, Takayuki; Sugiyama, Yasuo; Morimoto, Shinji

PA Takeda Chemical Industries Ltd, Japan

SO Jpn. Kokai Tokkyo Koho, 27 pp.

CODEN: JKXXAF

DT Patent

LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	JP 08295667	A2	19961112	JP 1995-129433	19950427

PRAI JP 1995-129433

19950427

OS MARPAT 126:89269

GI For diagram(s), see printed CA Issue.

AB The title compds. [I; A, B = (un)substituted (hetero)cycle; X = N, CR1; R, R1 = H, (un)substituted hydrocarbonyl; Y = (oxo)alkylene; Z = bond, alkylene; W = (un)substituted (hetero)cycle; when A, B = benzene ring, X = CR1, Y = CO, W = substituted cycle or (un)substituted heterocycle] are prepared I having a potent antagonism on tachykinin receptor (substance P receptor special) are useful as cholesterol acyltransferase (ACAT) inhibitors. Thus, N-[3,5-bis(trifluoromethyl)benzyl]-N'-(4-chloro-2-phenylaminophenyl)-N-methyloxamide (preparation given) was treated with HCl and reacted with AcONa in the presence of Pd/C under H atmospheric to give the title


compound (II). II showed IC50 of 0.36 nM against tachykinin receptors.

IT 185332-19-8P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(preparation of heterocyclic compds. as cholesterol acyltransferase inhibitors)

RN 185332-19-8 CAPLUS

CN 1H-Benzimidazole-2-acetamide, N-[2,6-bis(1-methylethyl)phenyl]-6-chloro-1-(2-pyridinyl)- (9CI) (CA INDEX NAME)



L30 ANSWER 6 OF 6 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1996:466914 CAPLUS

DN 125:142559

TI 4-Heterocyclylpiperidines promote release of growth hormone

IN Nargund, Ravi; Patchett, Arthur A.; Yang, Lihu

PA Merck and Co., Inc., USA

SO PCT Int. Appl., 84 pp.

CODEN: PIXXD2


DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9613265	A1	19960509	WO 1995-US13584	19951023
	W: AL, AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, IS, JP, KG, KR, KZ, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TT, UA, US, UZ				
	RW: KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
	US 5767118	A	19980616	US 1994-329357	19941026
	CA 2202784	AA	19960509	CA 1995-2202784	19951023
	AU 9539647	A1	19960523	AU 1995-39647	19951023
	EP 785784	A1	19970730	EP 1995-937576	19951023
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, NL, PT, SE				
	JP 10506914	T2	19980707	JP 1995-514657	19951023
PRAI	US 1994-329357	A1	19941026		
	WO 1995-US13584	W	19951023		
OS	MARPAT 125:142559				

GI



AB The present invention is directed to certain novel compds. identified as 4-heterocycle substituted piperidines I (R = benzimidazolyl, benzoxazinyl, pyridiyl, quinazoliny, etc., R1 = 3-phenylpropyl, benzyloxymethyl, indolylmethyl). These compds. promote the release of growth hormone in humans and animals. This property can be utilized to promote the growth of food animals to render the production of edible meat products more efficient, and in humans, to treat physiol. or medical conditions characterized by a deficiency in growth hormone secretion, such as short stature in growth hormone deficient children, and to treat medical conditions which are improved by the anabolic effects of growth hormone.

IT 179323-96-7P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)

(preparation of heterocyclylpiperidine growth hormone release promoters)

RN 179323-96-7 CAPLUS

CN 1H-Benzimidazole-2-propanoic acid, 1-[1-[2-[(2-amino-2-methyl-1-oxopropyl)amino]-3-(1H-indol-3-yl)-1-oxopropyl]-4-piperidinyl]-, ethyl ester, (R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

